

Nonperturbative QCD contribution to gluon-gluon- η' vertex

Mohammad R. Ahmady^{a*}, Victor Elias^{b†} and Emi Kou^{c‡}

^a*LINAC Laboratory, The Institute of Physical and Chemical Research (RIKEN)*

2-1 Hirosawa, Wako, Saitama 351-01, Japan

^b*Department of Applied Mathematics, University of Western Ontario*

London, Ontario, Canada N6A 5B7

^c*Department of Physics, Ochanomizu University*

1-1 Otsuka 2, Bunkyo-ku, Tokyo 112, Japan

(January 1998)

Abstract

We investigate the momentum dependence of the $g^* - g - \eta'$ vertex form factor in the presence of the nonperturbative QCD vacuum condensate. The behaviour of this form factor is crucial for, among other things, some proposed mechanisms devised to explain the large branching ratio for fast η' production in charmless B decays observed by CLEO. We show that the leading momentum dependence of this vertex form factor is not altered when the contribution of nonperturbative quark condensate is considered.

*Email: ahmady@riken.go.jp

†Email: zohar@apmaths.uwo.ca

‡Email: kou@fs.cc.ocha.ac.jp

The recent observation at CLEO of a large branching ratio for $B \rightarrow X_s \eta'$ decay [1] has led to various attempts for a theoretical explanation. Most of these attempts rely on η' being produced via its anomalous coupling to two gluons, which is parametrized by the form factor $H(q^2, p^2, m_\eta^2)$ with q and p being the four momenta of the gluons [2–4]. The momentum behaviour of this form factor is crucial for the success or failure of these proposed mechanisms. For example, $H(q^2, 0, m_{\eta'}^2) \approx H(0, 0, m_{\eta'}^2)$ has been assumed in Ref [2], so as to obtain results which agree with experiment. It has been pointed out, however, that a sigma-model calculation of $g^* - g - \eta'$ vertex (g^* and g are off-shell and real gluons, respectively) leads to a significant momentum dependence of the form $H(q^2, 0, m_{\eta'}^2) \sim 1/(q^2 - m_{\eta'}^2)$ [5]. In attempting to account for the $B \rightarrow X_s \eta'$ branching ratio, it is of interest to determine whether this leading momentum dependence is altered by nonperturbative QCD contributions. In this paper, we calculate the quark condensate component of the nonperturbative contribution to H to investigate whether the insertion of the $\langle \bar{f}f \rangle$ augmented quark propagator in the triangle loop alters the leading momentum dependence of the vertex form factor. The purely-perturbative form factor $H(q^2, 0, m_{\eta'}^2)$ for the vertex $g^* - g - \eta'$ (Fig. 1) is found to be

$$V_{\mu\nu} = \sum_{f=u,d,s} \frac{g_f^{\eta'} g_s^2 m_f}{2\pi^2} H(q^2, 0, m_{\eta'}^2) \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta, \quad (1)$$

with

$$H(q^2, 0, m_{\eta'}^2) = \frac{1}{q^2 - m_{\eta'}^2} \left[I\left(\frac{m_{\eta'}^2}{m_f^2}\right) - I\left(\frac{q^2}{m_f^2}\right) \right], \quad (2)$$

$$I(x^2) = \begin{cases} -2\arcsin^2 \frac{|x|}{2} & 0 \leq x^2 \leq 4, \\ 2 \left[\text{Ln}\left(\frac{|x|}{2} + \sqrt{\frac{x^2}{4} - 1}\right) - \frac{i\pi}{2} \right]^2 & x^2 \geq 4. \end{cases}$$

$g_f^{\eta'}$ is the coupling of η' to the quark of flavour f , and g_s is the strong interaction coupling constant. We note from Eq. (2) that the leading momentum dependence of $H \sim 1/(q^2 - m_{\eta'}^2)$ indicates a significant suppression of this form factor at large q^2 . As a result, the mechanism suggested in Ref. [2] for $B \rightarrow X_s \eta'$ decay falls short of experimental data by at least an order

of magnitude. A reconciliation of this mechanism to experiment, however, may be possible from a hardening of the leading momentum dependence of H by the nonperturbative content of QCD, content which is not taken into account in Eq. (2). We investigate this possibility by calculating the quark-condensate contributions to the form factor H . QCD condensates characterize the nonperturbative content of the QCD vacuum, and therefore, the inclusion of a quark-condensate insertion should give us some indication of the leading nonperturbative contributions to H . We proceed first by replacing the usual perturbative quark propagator in Fig. 1 with the full quark propagator $S(k)$ [6]:

$$S(k) = S^{\text{P}}(k) + S^{\text{NP}}(k) = -i \int d^4x e^{ik \cdot x} \langle \Omega | T \Psi(x) \bar{\Psi}(0) | \Omega \rangle , \quad (3)$$

where $|\Omega \rangle$ is the nonperturbative QCD vacuum. As a result, the nonperturbative contributions to the quark propagator can be written as

$$iS^{\text{NP}}(k) = \int d^4x e^{ik \cdot x} \langle \Omega | : \Psi(x) \bar{\Psi}(0) : | \Omega \rangle . \quad (4)$$

The nonlocal vacuum expectation value (vev) in Eq. (4) can be expanded in terms of local condensates [7]. To ascertain how leading nonperturbative QCD contributions affect the form factor H , only the lowest-dimensional (quark-condensate) component of the expansion is considered below. The quark-condensate projection of the nonperturbative quark propagator is taken from Ref. [6],

$$iS^{\text{NP}}(k) = (2\pi)^4 (\gamma_\mu k^\mu + m_f) F(k), \quad (5)$$

where the Fourier transform of $F(k)$ is given by

$$\int d^4k e^{-ik \cdot x} F(k) = -\frac{\langle \bar{f} f \rangle}{6m_f^2} \frac{J_1(m_f \sqrt{x^2})}{\sqrt{x^2}} . \quad (6)$$

$F(k)$ has the following important on-shell property [6]:

$$(k^2 - m_f^2) F(k) = 0 . \quad (7)$$

The dimension-3 quark condensate $\langle \bar{f} f \rangle$ is the vev of the normal ordered local two-quark fields, i.e.

$$\langle \bar{f}f \rangle = \langle \Omega | : \bar{\Psi}(0)\Psi(0) : | \Omega \rangle . \quad (8)$$

Using the Feynman rule of the Eq. (5), one can obtain the nonperturbative $\langle \bar{f}f \rangle$ contribution to H in a straightforward manner. The relevant Feynman diagrams are illustrated in Fig. 2 where the nonperturbative quark propagator S^{NP} is depicted by a disconnected line with two dots. For example, the expression for Fig 2a is written as follows:

$$\begin{aligned} {}^{(2a)}V_{\mu\nu}^{\langle \bar{f}f \rangle} &= -4g_f^{\eta'} g_s^2 m_f \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta \\ &\times \int d^4k \int_0^1 dz \frac{F(k)}{[-zq^2 - 2k \cdot \{z(q+p) - p\}]^2} , \end{aligned} \quad (9)$$

where the gluon with four-momentum p is taken to be on-shell. To proceed, we use the following formulae:

$$\begin{aligned} \int d^4k \frac{F(k)}{[a - k \cdot v + i\epsilon]^2} &= - \int d^4k F(k) \int_0^\infty d\eta \eta e^{i\eta(a - k \cdot v + i\epsilon)} , \\ &= \frac{\langle \bar{f}f \rangle}{6m_f^2} \int_0^\infty d\eta e^{i\eta a} \frac{J_1(m_f \eta \sqrt{v^2})}{\sqrt{v^2}} , \\ &= \frac{\langle \bar{f}f \rangle}{6m_f^3 v^2} \left(1 - \frac{1}{\sqrt{1 - \frac{m_f^2 v^2}{a^2}}} \right) , \end{aligned} \quad (10)$$

where the second line is obtained from Eq. (6), and the final line from differentiation (with respect to a) of a tabulated integral [8], corresponding to the second line of (10). Consequently, the contribution of Fig 2a (Eq. (9)) is

$$\begin{aligned} {}^{(2a)}V_{\mu\nu}^{\langle \bar{f}f \rangle} &= -\frac{g_f^{\eta'} g_s^2 \langle \bar{f}f \rangle}{6m_f^2} \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta \\ &\times \int_0^1 dz \frac{1}{z(zm_{\eta'}^2 + q^2 - m_{\eta'}^2)} \left[1 - \frac{1}{\sqrt{1 - \frac{4m_f^2(zm_{\eta'}^2 + q^2 - m_{\eta'}^2)}{zq^4}}} \right] . \end{aligned} \quad (11)$$

Expressions for Figs. 2b and 2c are calculated in the same way, leading to the following aggregate quark-condensate contribution to $g^* - g - \eta'$ vertex:

$$V_{\mu\nu}^{\langle \bar{f}f \rangle} = \sum_{f=u,d,s} \frac{g_f^{\eta'} g_s^2 m_f}{2\pi^2} H^{\langle \bar{f}f \rangle}(q^2, 0, m_{\eta'}^2) \epsilon_{\mu\nu\alpha\beta} q^\alpha p^\beta , \quad (12)$$

with

$$H^{<\bar{f}f>}(q^2, 0, m_{\eta'}^2) = -\frac{\pi^2 <\bar{f}f>}{3m_f^3} \frac{1}{q^2 - m_{\eta'}^2} G\left(\frac{q^2}{m_{\eta'}^2}, \frac{m_f^2}{m_{\eta'}^2}\right) , \quad (13)$$

where

$$G(x, y) = \text{Ln} \left[\frac{4y^3 \left[(x-1)(\sqrt{x(x-4y)} + 2y) - x^2 + x(x-1)^2 \right]}{[2y(x-1) + x^3 - 2x^2] \left[(2y-1)\sqrt{1-4y} + 2y^2 - 4y + 1 \right] \left[\sqrt{x(x-4y)} + x - 2y \right]} \right] .$$

We note that the leading momentum dependence of the quark condensate contribution to the vertex form factor is indeed the same as in (2); *i.e.*, $H^{<\bar{f}f>} \sim 1/(q^2 - m_{\eta'}^2)$. Therefore, we conclude that the lowest-dimensional contribution from nonperturbative QCD *does not* alter the significant suppression of the $g^* - g - \eta'$ vertex for large gluon virtualities. Whether or not this result is valid when higher dimensional condensates are considered remains to be investigated. However, one would anticipate the effect of higher-dimensional order parameters to be negligible at large momenta, suggesting that the momentum-dependence of $g^* - g - \eta'$ vertex form factor is not altered appreciably by nonperturbative QCD effects.

ACKNOWLEDGEMENTS

M. A. acknowledges support from the Science and Technology Agency of Japan under an STA fellowship. V.E. acknowledges support from the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- [1] Jim Smith, talk given at the 7th International Symposium on Heavy Flavor Physics, Santa Barbara CA (July 1997); B. Behren, talk given at the Princeton BaBar meeting (March 1997); P. Kim, talk given at FCNC 97, Santa Monica CA (Feb,1997); F. Würthwein, CALT-68-2121, hep-ex/9706010.
- [2] D. Atwood and A. Soni, Phys. Lett. **B405** 150 (1997).
- [3] W.-S. Hou and B. Tseng, hep-ph/9705304.
- [4] M. R. Ahmady, E. Kou and A. Sugamoto, hep-ph/9710509.
- [5] A. L. Kagan and A. A. Petrov, hep-ph/9707354.
- [6] E. Bagan, M. R. Ahmady, V. Elias and T. G. Steele, Z. Phys. **C61** 157 (1994).
- [7] V. Elias, T. G. Steele and M. D. Scadron, Phys. Rev. **D38**, 1584 (1988).
- [8] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products* (Academic Press, Orlando, 1980) p 712, Eq (6.623.3).

Figure Captions

Figure 1: Quark loop triangle diagram for $g - g - \eta'$ vertex.

Figure 2: Nonperturbative quark condensate contributions to $g - g - \eta'$ vertex.

Fig. 1




Fig. 2

